# 

Please write clearly in	n block capitals.	
Centre number	Candidate number	
Surname		
Forename(s)		
Candidate signature	I declare this is my own work.	,

## GCSE COMBINED SCIENCE: TRILOGY

Higher Tier Chemistry Paper 1H

### Time allowed: 1 hour 15 minutes

#### Materials

For this paper you must have:

- a ruler
- a scientific calculator
- the periodic table (enclosed).

#### Instructions

- Use black ink or black ball-point pen.
- · Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer all questions in the spaces provided.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.
- In all calculations, show clearly how you work out your answer.

#### Information

- The maximum mark for this paper is 70.
- The marks for questions are shown in brackets.
- You are expected to use a calculator where appropriate.
- You are reminded of the need for good English and clear presentation in your answers.



For Examiner's Use					
Question	Mark				
1					
2					
3					
4					
5					
6					
7					
TOTAL					



This question is about the periodic table.

**1 Figure 1** shows part of Mendeleev's version of the periodic table.

Figure 1	
----------	--

н																	
Li	i	В	е		В			С			N		C		F		
Na	а	М	g		Al			Si			Ρ		S		Cl		
К	Cu	Са	Zn				Ti			V	А	s	Cr	Se	Mn	Br	Fe Co Ni
Rb	Ag	Sr	Cd	Y		In	Zr		Sn	Nb	S	b	Мо	Те		I	Ru Rh Pd

Which group of elements had **not** been discovered when Mendeleev's version of the periodic table was published?

[1 mark]



0 1

0 1





Turn over ►

4

	Potassium has dif	ferent isotopes.			Do not write outside the box
0 1.4	What is meant by				
	You should refer to	o subatomic particles		[2 marks]	
0 1.5	Table 1 shows theisotopes of potass		the percentage abundance of	<sup>f</sup> two	
			Table 1		
		Mass number	Percentage abundance		
		39	93.1		
		41	6.9		
	Calculate the relat	ive atomic mass (A <sub>r</sub> )	of potassium.		
		to 1 decimal place.			
				[3 marks]	
		Relative a	tomic mass (1 decimal place)	) =	8







0 2	Acids react to produce salts.	Do not write outside the box
	Universal indicator is added to water and then nitric acid is added to the mixture.	
02.1	Give the colour change when nitric acid is added to the mixture of universal indicator and water. [1 mark]	
	Tick (✓) <b>one</b> box.	
	Blue to red	
	Green to purple	
	Green to red	
	Red to purple	
02.2	What happens to the pH of water when nitric acid is added? [1 mark]	
	Tick (✓) <b>one</b> box.	
	Decreases	
	Stays the same	
	Increases	
02.3	What is the state symbol for nitric acid?	
	[1 mark]	



		Do not write
	Zinc carbonate reacts with nitric acid.	outside the box
	The word equation for the reaction is:	
	zinc carbonate + nitric acid $\rightarrow$ zinc nitrate + water + carbon dioxide white solid colourless solution	
02.4	Give <b>two</b> observations that would be made when zinc carbonate is added to nitric acid until the zinc carbonate is in excess. [2 marks]	
	1	
	2	
	The formula of the zinc ion is Zn <sup>2+</sup>	
0 2 . 5	The formula of the nitrate ion is $NO_3^-$	
	What is the formula for zinc nitrate? [1 mark]	
	Tick (✓) <b>one</b> box.	
	ZnNO <sub>3</sub>	
	Zn(NO <sub>3</sub> ) <sub>2</sub>	
	Zn <sub>2</sub> NO <sub>3</sub>	
	Zn <sub>2</sub> (NO <sub>3</sub> ) <sub>2</sub>	
	Question 2 continues on the next page	



Turn over ►

02.6	Acids react with insoluble metal oxides to produce salts.	outside the box
	Plan a method to produce a pure, dry sample of the soluble salt copper chloride from an acid and a metal oxide.	
	[6 marks]	
		12



Do not write













Turn over ►









0 4	Carbon can exist in a number of different structures.	Do not write outside the box
04.1	The first fullerene to be discovered was Buckminsterfullerene?       [1 mark]         Tick (~) one box.       [1 mark]         C <sub>40</sub>	
04.2	Graphite is a form of carbon. Explain why graphite conducts electricity. [2 marks]	



	Steel is an alloy	of iron and ca	arbon.				Do not wri outside th box		
04.3	Explain why stee	el is harder tha	an iron.			[3 marks]	1		
							_		
							_		
							_		
							-		
04.4	Iron is alloyed w	ith carbon and	d other metal	s to make st	ainless steel.				
	A stainless steel								
	Table 2 shows t	he mass of ea	ach element i	n the fork.					
	Table 2								
	Element		Iron	Carbon	Chromium	Nickel			
	Mass of el	ement in g	X	0.05	10.44	5.80			
	Calculate the ma	ass of iron ( <b>X</b> )	in the fork.						
		. ,				[4 marks	]		
							_		
							_		
							_		
							=		
							-		
					X = _	g			



0 5	This question is about the electrolysis of aqueous solutions.	Do not write outside the box
	Hydrogen gas and chlorine gas are produced when sodium chloride solution is electrolysed.	
0 5.1	Hydrogen ions (H <sup>+</sup> ) are attracted to the negative electrode.	
	The half equation for the reaction at the negative electrode is:	
	$2  H^+$ + $2  e^- \rightarrow H_2$	
	What type of reaction happens at the negative electrode?	
	Give the reason for your answer. [2 marks]	
	Type of reaction	
	Reason	
	Chloride ions are attracted to the positive electrode.	
0 5 . 2	Complete the half equation for the production of chlorine gas ( $Cl_2$ ).	
	[2 marks]	
	$Cl^- \rightarrow +$	



0 5.3	Hydrogen gas and oxygen gas are produced when sodium sulfate solution is electrolysed.	Do not write outside the box
	Explain how oxygen gas is produced in the electrolysis of sodium sulfate solution. [4 marks]	
		8
	Turn over for the next question	
	Turn over ►	

IB/M/Jun21/8464/C/1H

06	Metal oxides are produced when metals are heated in air.	Do not write outside the box
	A student investigated the change in mass when 0.12 g of magnesium was heated in air.	
	Figure 5 shows the apparatus.	
	Figure 5	
	Lid	
	Crucible	
	Tripod Heat Balance	
	The student measured the mass of magnesium oxide produced.	
06.1	0.12 g of magnesium reacted to produce 0.20 g of magnesium oxide.	
	Calculate the number of moles of oxygen gas (O <sub>2</sub> ) that reacted.	
	Relative atomic mass ( $A_r$ ): O = 16	
	[3 marks]	
	Moles of oxygen gas =	



06.2	The student repeated the experiment <b>without</b> a lid on the crucible.	Do not write outside the box
	Suggest why the mass of magnesium oxide produced would be different without a lid on the crucible.	
	[2 marks]	
0 6 . 3	Copper reacts with oxygen to produce copper oxide.	
	63.5 g of copper produces 79.5 g of copper oxide.	
	Calculate the mass of copper oxide produced when 0.50 g of copper reacts with oxygen.	
	Give your answer to 3 significant figures. [3 marks]	
	Mass (3 significant figures) =g	
	Question 6 continues on the next page	

		Do not write
0 6.4	Iron reacts with oxygen to produce an oxide of iron.	outside the box
	0.015 moles of iron reacts with 0.010 moles of oxygen gas $(O_2)$ .	
	Determine:	
	<ul> <li>the formula of the iron oxide produced</li> </ul>	
	<ul> <li>the balanced symbol equation for the reaction.</li> </ul>	
	[4 marks]	
	Formula of iron oxide =	
	Balanced symbol equation	
		12
	·	







0 7	Methane and wate	e, ethane, propane and butane er.	all react with	oxygen to pr	oduce carboi	n dioxide
0 7.1		why a mixture of methane and in terms of particles.	l oxygen does	s <b>not</b> react at		
					[4	2 marks]
07.2		shows the energy released wh gen to produce carbon dioxide		ethane and p	propane reac	t
			Compoun	d reacted w	ith oxygen	_
			Methane	Ethane	Propane	-
		Formula of compound	CH <sub>4</sub>	$C_2H_6$	C <sub>3</sub> H <sub>8</sub>	_
		Energy released in kJ/mol	680	1160	1640	
		he energy released when buta lioxide and water.	ne (C₄H₁₀) rea	acts with oxy	-	ce [1 mark]
			Energy	released = _		_kJ/mol



**0 7**. **3** Propane reacts with oxygen to produce carbon dioxide and water.

The displayed formula equation for the reaction is:

$$\begin{array}{ccccccc} H & H & H \\ | & | & | \\ H - C - C - C - C - H & + 5 & 0 = 0 & \longrightarrow & 3 & 0 = C = 0 & + 4 & H - 0 - H \\ | & | & | \\ H & H & H \end{array}$$

The reaction is exothermic.

In the reaction, the energy released when forming new bonds is 1640 kJ/mol greater than the energy needed when breaking bonds.

Table 4 shows bond energies.

Table 4

Bond	H–C	C–C	0=0	C=0	0–Н
Bond energy in kJ/mol	410	x	500	740	460

END OF QUESTIONS

Calculate the C—C bond energy (X).

[5 marks]

kJ/mol

X =



Do not write outside the box





Question number	Additional page, if required. Write the question numbers in the left-hand margin.	Do not write outside the box



Question number	Additional page, if required. Write the question numbers in the left-hand margin.



Question number	Additional page, if required. Write the question numbers in the left-hand margin.



Question number	Additional page, if required. Write the question numbers in the left-hand margin.
	Copyright information
	For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from www.aqa.org.uk.
	Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.
	Copyright © 2021 AQA and its licensors. All rights reserved.



