

Section 3: Surds

Solutions to Exercise

$$1. \text{ (i)} \quad \sqrt{8} = \sqrt{4 \times 2} = \sqrt{4} \times \sqrt{2} = 2\sqrt{2}$$

$$\text{(ii)} \quad \sqrt{50} = \sqrt{25 \times 2} = \sqrt{25} \times \sqrt{2} = 5\sqrt{2}$$

$$\text{(iii)} \quad \sqrt{48} = \sqrt{16 \times 3} = \sqrt{16} \times \sqrt{3} = 4\sqrt{3}$$

$$\text{(iv)} \quad \sqrt{216} = \sqrt{36 \times 6} = \sqrt{36} \times \sqrt{6} = 6\sqrt{6}$$

$$\text{(v)} \quad \sqrt{63} = \sqrt{9 \times 7} = \sqrt{9} \times \sqrt{7} = 3\sqrt{7}$$

$$\text{(vi)} \quad \sqrt{300} = \sqrt{100 \times 3} = \sqrt{100} \times \sqrt{3} = 10\sqrt{3}$$

$$\text{(vii)} \quad \sqrt{6} \times \sqrt{27} = \sqrt{3} \times \sqrt{2} \times \sqrt{9} \times \sqrt{3} = \sqrt{2} \times 3 \times 3 = 9\sqrt{2}$$

$$\text{(viii)} \quad \sqrt{12} \times \sqrt{15} = \sqrt{4} \times \sqrt{3} \times \sqrt{3} \times \sqrt{5} = 2 \times 3 \times \sqrt{5} = 6\sqrt{5}$$

$$\begin{aligned} \text{(ix)} \quad \sqrt{10} \times \sqrt{24} \times \sqrt{15} &= \sqrt{5} \times \sqrt{2} \times \sqrt{4} \times \sqrt{3} \times \sqrt{2} \times \sqrt{3} \times \sqrt{5} \\ &= 5 \times 2 \times 2 \times 3 \\ &= 60 \end{aligned}$$

$$2. \text{ (i)} \quad (1 + \sqrt{2}) + (3 - 2\sqrt{2}) = 1 + 3 + \sqrt{2} - 2\sqrt{2} \\ = 4 - \sqrt{2}$$

$$\text{(ii)} \quad (5\sqrt{2} - 2\sqrt{3}) - (\sqrt{2} + 3\sqrt{3}) = 5\sqrt{2} - 2\sqrt{3} - \sqrt{2} - 3\sqrt{3} \\ = 4\sqrt{2} - 5\sqrt{3}$$

$$\text{(iii)} \quad 2(\sqrt{5} - 3\sqrt{3}) + 3(2\sqrt{5} + \sqrt{3}) = 2\sqrt{5} - 6\sqrt{3} + 6\sqrt{5} + 3\sqrt{3} \\ = 8\sqrt{5} - 3\sqrt{3}$$

$$\text{(iv)} \quad \sqrt{18} + \sqrt{72} - \sqrt{98} = \sqrt{9 \times 2} + \sqrt{36 \times 2} - \sqrt{49 \times 2} \\ = 3\sqrt{2} + 6\sqrt{2} - 7\sqrt{2} \\ = 2\sqrt{2}$$

AQA FM Number and Algebra 1 Exercise solutions

3. (i) $(1 + \sqrt{2})(3 - \sqrt{2}) = 3 - \sqrt{2} + 3\sqrt{2} - 2$
 $= 1 + 2\sqrt{2}$

(ii) $(2 - \sqrt{3})(3 + 2\sqrt{3}) = 6 + 4\sqrt{3} - 3\sqrt{3} - 2 \times 3$
 $= \sqrt{3}$

(iii) $(3 - 2\sqrt{5})(1 - 3\sqrt{5}) = 3 - 9\sqrt{5} - 2\sqrt{5} + 6 \times 5$
 $= 33 - 11\sqrt{5}$

(iv) $(\sqrt{2} + 2\sqrt{3})(5\sqrt{2} - \sqrt{3}) = 5 \times 2 - \sqrt{2}\sqrt{3} + 10\sqrt{3}\sqrt{2} - 2 \times 3$
 $= 10 - \sqrt{6} + 10\sqrt{6} - 6$
 $= 4 + 9\sqrt{6}$

(v) $(\sqrt{7} + \sqrt{2})(\sqrt{7} - \sqrt{2}) = 7 - \sqrt{7}\sqrt{2} + \sqrt{2}\sqrt{7} - 2$
 $= 5$

(vi) $(3 - \sqrt{2})^2 = (3 - \sqrt{2})(3 - \sqrt{2})$
 $= 9 - 3\sqrt{2} - 3\sqrt{2} + 2$
 $= 11 - 6\sqrt{2}$

4. (i) $\frac{3}{\sqrt{3}} \times \frac{\sqrt{3}}{\sqrt{3}} = \frac{3\sqrt{3}}{3} = \sqrt{3}$

(ii) $\frac{1}{\sqrt{5}} \times \frac{\sqrt{5}}{\sqrt{5}} = \frac{\sqrt{5}}{5}$

(iii) $\frac{1 + \sqrt{2}}{\sqrt{2}} \times \frac{\sqrt{2}}{\sqrt{2}} = \frac{(1 + \sqrt{2})\sqrt{2}}{2} = \frac{\sqrt{2} + 2}{2}$

(iv) $\frac{1}{\sqrt{3} + 1} \times \frac{\sqrt{3} - 1}{\sqrt{3} - 1} = \frac{\sqrt{3} - 1}{(\sqrt{3} + 1)(\sqrt{3} - 1)} = \frac{\sqrt{3} - 1}{3 - 1} = \frac{\sqrt{3} - 1}{2}$

(v) $\frac{\sqrt{2}}{2 - \sqrt{2}} \times \frac{2 + \sqrt{2}}{2 + \sqrt{2}} = \frac{\sqrt{2}(2 + \sqrt{2})}{(2 - \sqrt{2})(2 + \sqrt{2})} = \frac{2\sqrt{2} + 2}{4 - 2} = \frac{2\sqrt{2} + 2}{2} = \sqrt{2} + 1$

AQA FM Number and Algebra 1 Exercise solutions

$$\begin{aligned}
 (\text{vi}) \quad & \frac{1-\sqrt{3}}{2-\sqrt{3}} \times \frac{2+\sqrt{3}}{2+\sqrt{3}} = \frac{(1-\sqrt{3})(2+\sqrt{3})}{(2-\sqrt{3})(2+\sqrt{3})} \\
 &= \frac{2+\sqrt{3}-2\sqrt{3}-3}{4-3} \\
 &= \frac{-1-\sqrt{3}}{1} \\
 &= -1-\sqrt{3}
 \end{aligned}$$

$$\begin{aligned}
 (\text{vii}) \quad & \frac{1+2\sqrt{5}}{3-\sqrt{5}} \times \frac{3+\sqrt{5}}{3+\sqrt{5}} = \frac{(1+2\sqrt{5})(3+\sqrt{5})}{(3-\sqrt{5})(3+\sqrt{5})} \\
 &= \frac{3+\sqrt{5}+6\sqrt{5}+2\times 5}{9-5} \\
 &= \frac{13+7\sqrt{5}}{4}
 \end{aligned}$$

$$\begin{aligned}
 (\text{viii}) \quad & \frac{1+\sqrt{2}}{\sqrt{3}+\sqrt{2}} \times \frac{\sqrt{3}-\sqrt{2}}{\sqrt{3}-\sqrt{2}} = \frac{(1+\sqrt{2})(\sqrt{3}-\sqrt{2})}{(\sqrt{3}+\sqrt{2})(\sqrt{3}-\sqrt{2})} \\
 &= \frac{\sqrt{3}-\sqrt{2}+\sqrt{6}-2}{3-2} \\
 &= \sqrt{3}-\sqrt{2}+\sqrt{6}-2
 \end{aligned}$$

$$\begin{aligned}
 (\text{ix}) \quad & \frac{\sqrt{6}+\sqrt{3}}{\sqrt{6}-\sqrt{3}} \times \frac{\sqrt{6}+\sqrt{3}}{\sqrt{6}+\sqrt{3}} = \frac{(\sqrt{6}+\sqrt{3})(\sqrt{6}+\sqrt{3})}{(\sqrt{6}-\sqrt{3})(\sqrt{6}+\sqrt{3})} \\
 &= \frac{6+\sqrt{18}+\sqrt{18}+3}{6-3} \\
 &= \frac{9+2\sqrt{18}}{3}
 \end{aligned}$$

5. One possible width is $(\sqrt{7}-1)$ cm.

In this case, the area is $(\sqrt{7}+1)(\sqrt{7}-1) = 7-1 = 6$ cm².

Other answers are possible.